



TITLE:

# THE Nil-Nil THEOREM IN ALGEBRAIC $K$ -THEORY (Geometry of Transformation Groups and Related Topics)

AUTHOR(S):

KHAN, QAYUM

---

CITATION:

KHAN, QAYUM. THE Nil-Nil THEOREM IN ALGEBRAIC  $K$ -THEORY (Geometry of Transformation Groups and Related Topics). 数理解析研究所講究録 2008, 1612: 99-100

ISSUE DATE:

2008-09

URL:

<http://hdl.handle.net/2433/140074>

RIGHT:

# THE Nil-Nil THEOREM IN ALGEBRAIC $K$ -THEORY

QAYUM KHAN

The reduced Nil-groups are certain reduced  $K$ -theory groups defined by Friedhelm Waldhausen for pure amalgams and tensor algebras [1]. They measure the defect in his Mayer–Vietoris sequence in the algebraic  $K$ -theory of rings. In the main paper [2], we recently showed that the apparently more complicated amalgam Nil can be computed in terms of the apparently simpler tensor Nil.

**Theorem 1.** *Let  $R$  be a (unital, associative) ring. Let  $\mathcal{B}_1$  and  $\mathcal{B}_2$  be  $R$ -bimodules. Suppose  $I$  is a small, filtered category and  $\mathcal{B}_2 = \operatorname{colim}_{\alpha \in I} \mathcal{B}_2^\alpha$  is a direct limit of  $R$ -bimodules such that the left  $R$ -module structure of each  $\mathcal{B}_2^\alpha$  is finitely generated and projective. Then, for every  $n \in \mathbb{Z}$ , there is an induced isomorphism*

$$\tilde{K}_n(j) : \tilde{\operatorname{Nil}}_n(R; \mathcal{B}_1, \mathcal{B}_2) \longrightarrow \tilde{\operatorname{Nil}}_n(R; \mathcal{B}_1 \otimes_R \mathcal{B}_2).$$

An important special case are those amalgams of group rings which are induced by an epimorphism onto the infinite dihedral group  $D_\infty = \mathbb{Z}/2 * \mathbb{Z}/2 = \mathbb{Z} \rtimes_{-1} \mathbb{Z}/2$ .

**Corollary 2.** *Suppose  $G$  is a group with an epimorphism  $p : G \rightarrow D_\infty$ . Denote the  $p$ -induced injective amalgamated product decomposition  $G = G_1 *_F G_2$ . Consider the index-two subgroup  $\bar{G} := p^{-1}(\mathbb{Z})$  of  $G$ . Denote the  $p$ -induced injective HNN-extension  $\bar{G} = F \rtimes_\alpha \mathbb{Z}$ . Then, for all rings  $R$  and for all  $n \in \mathbb{Z}$ , there is an isomorphism of abelian groups:*

$$\tilde{\operatorname{Nil}}_n(R[F]; R[G_1 - F], R[G_2 - F]) \cong NK_{n+1}(R[F], \alpha).$$

The right-hand side of the isomorphism is the twisted Bass Nil-group [3] of F.T. Farrell and W.C. Hsiang [4]. These are more readily computable since they involve the Wang sequence in  $K$ -theory of the twisted polynomial ring  $R[F]_\alpha[x]$ .

The following application [2] of the above corollary is a sharpening of the fibered isomorphism conjecture of F. T. Farrell and L. E. Jones in algebraic  $K$ -theory. Given a group  $G$ , denote  $\operatorname{vc}$  as the class of virtually cyclic subgroups and  $\operatorname{fbc}$  as the subclass of finite-by-cyclic subgroups. The elements of the complement  $\operatorname{vc} - \operatorname{fbc}$  are exactly those subgroups of  $G$  which are finite-by- $D_\infty$ .

**Theorem 3.** *Let  $\varphi : \Gamma \rightarrow G$  be an epimorphism of groups. Then, for all rings  $R$  and for all  $n \in \mathbb{Z}$ , the following induced map is an isomorphism:*

$$H_n^\Gamma(E_{\varphi^* \operatorname{fbc}} \Gamma; \mathbf{K}_R) \longrightarrow H_n^\Gamma(E_{\varphi^* \operatorname{vc}} \Gamma; \mathbf{K}_R).$$

Both sides are equivariant homology groups, whose coefficients are given by the spectrum-valued functor  $\mathbf{K}_R : \operatorname{Or} \Gamma \rightarrow \operatorname{SPECTRA}$  of the Bredon orbit category [5].

Recently, the Farrell–Jones conjecture has been proven by A. Bartels, W. Lück, and H. Reich for a large class of infinite groups with torsion [6]. Consider the non-fibered case of  $\varphi = \operatorname{id}_\Gamma$ .

**Corollary 4.** *Let  $\Gamma$  be a word-hyperbolic group. Then, for all rings  $R$  and for all  $n \in \mathbb{Z}$ , the algebraic  $K$ -theory assembly map is an isomorphism:*

$$H_n^\Gamma(E_{\text{fbc}}\Gamma; \mathbf{K}_R) \longrightarrow K_n(R[\Gamma]) .$$

This isomorphism yields specific fruit. In the following calculation [2], the Bass  $NK$ -groups vanish if  $R$  is a regular Noetherian ring, such as if  $R = \mathbb{Z}$ . Here  $K_n(R[x]) = K_n(R) \oplus NK_n(R)$  by definition.

**Theorem 5.** *Consider the modular group  $\Gamma = PSL(2, \mathbb{Z})$ . Then, for any ring  $R$  and integer  $n$ , we have*

$$K_n(R[\Gamma]) = (K_n(R[\mathbb{Z}/2]) \oplus K_n(R[\mathbb{Z}/3])) / K_n(R) \oplus \bigoplus_{\mathbb{N}_0} NK_n(R) .$$

Finally, [2] provides the first example of a non-vanishing amalgam Nil-group.

*Example 6.* Consider the group  $G_0 := \mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}$ . Then

$$\widetilde{\text{Nil}}_0(\mathbb{Z}[G_0]; \mathbb{Z}[G_0], \mathbb{Z}[G_0]) = NK_1(\mathbb{Z}[G_0])$$

is a non-zero abelian group (see [3]), which is a summand of the Whitehead group  $\text{Wh}(G_0 \times D_\infty)$ . Therefore we obtain the following topological consequence. Consider the finite CW-complexes

$$\begin{aligned} W &:= \mathbb{RP}^2 \times \mathbb{RP}^2 \times S^1 \\ X &:= W \times S^2 \\ Y &:= W \times (\mathbb{RP}^3 - \text{int } D^3) . \end{aligned}$$

Given any non-zero element of the above amalgam Nil-group, one can construct [7] the first known example of a homotopy equivalence  $h : K \rightarrow Y \cup_X Y$ , where  $K$  is a certain finite CW-complex, such that  $h$  is **not** splittable along  $X$ . Here, we say  $h$  is *splittable along  $X$*  if there exist a simple homotopy equivalence  $s : K' \rightarrow K$  of finite CW-complexes and a homotopy equivalence  $h' : K' \rightarrow Y \cup_X Y$  such that  $h \circ s \simeq h'$  and the cellular restriction  $(h')^{-1}(X) \rightarrow X$  is a homotopy equivalence.

## REFERENCES

- [1] Friedhelm Waldhausen. Algebraic  $K$ -theory of generalized free products. *Ann. of Math. (2)*, 108(1):135–256, 1978.
- [2] James F. Davis, Qayum Khan, Andrew Ranicki. Algebraic  $K$ -theory over the infinite dihedral group, submitted. URL <http://arxiv.org/abs/0803.1639>
- [3] Hyman Bass. Algebraic  $K$ -theory. W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [4] F. T. Farrell and W. C. Hsiang. Manifolds with  $\pi_1 = G \rtimes_\alpha T$ . *Amer. J. Math.*, 95:813–848, 1973.
- [5] James F. Davis and Wolfgang Lück. Spaces over a category and assembly maps in isomorphism conjectures in  $K$ - and  $L$ -theory. *K-Theory*, 15(3):201–252, 1998.
- [6] Arthur Bartels, Wolfgang Lück, and Holger Reich. The  $K$ -theoretic Farrell-Jones Conjecture for hyperbolic groups. *Invent. Math.*, 172:29–70, 2008.
- [7] Friedhelm Waldhausen. Whitehead groups of generalized free products. *Unpublished paper with erratum*, 1969. URL <http://www.maths.ed.ac.uk/~aar/surgery/whgen.pdf>

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE TN 37240 U.S.A.  
E-mail address: qayum.khan@vanderbilt.edu